Missouri Botanical Garden

Peter H. Raven Library

George Engelmann Botanical Notebooks



Pagination Note:
Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.

Cuscuteae for Torrey & Gray's flora

1/

December 1846

Convolvulaceae

suborder Cuscuteae Chois. conv.

Lobes of calyx united or rarely distinct; stamens equal in number, and alternating with the divisions of corolla adhering to the tube of the corolla, rarely naked generally bearing near their base tufts or wing like toothed or fringed appendages, which mostly are connected at their free upper part mostly are united into one scale; these scales opposite the filaments, are again frequently connected at the base into one

Lobes of corolla imbricate in aestivation; ovary two celled; embryo elongated, filiform, without cotyledons, spirally coiled around a fleshy, mucilaginous albumen.

Annual Parasitics, with filiform white, yellow or red stems and scale-like leaves, without verdure, twining to the left, germinating in the ground, but soon fastening on (by their papillose roots, which penetrate the epidermis of) those plants from which they draw their sustenance mostly dicotyledonous ones, innocuous or poisonous as Euphorbia, Rhus Toxicodendron &c, rarely monocotyledonous, never cryptogamous plants — (though sometimes their limbs root) generally herbaceous, but also shrubby, and (big) trees) and then loosing their connection with the soil.

Calyx persistent, its lobes united or rarely distinct; corolla campanulate globular or tubular, with 5 or rarely 4 lobes; stamens equal in number and alternating with the lobes of the corolla, filaments adhering to its tube; rarely naked, generally bearing on each side near the base winglike tooth appendages which at their free upper end are mostly united into one toothed or fringed, scale; ovary two celled, with 4 erect ovules, styles 2, distinct or united, stigmas elongated or capitate, (surrounded at base or covered by the bitten corolla with the stamens) capsule 2 celled, 4 valved (frequently by abortion less) seeds either opening by circumscission, when the base with part or the whole or not opening at all. in either case the two carpels of the capsule of those, which have two distinct styles

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

separate to some extent at the top, leaving an opening between the styles, which however does not penetrate into the capsule, but only between two lamellae of the dissepiment. — Seeds large, yellowish compressed & irregularly ovoid, with a rough surface, yellowish or brown.

1. Cuscuta Linn. partly ex parte / Cuscuta and Epilinella Pfeiff. Styles two, distinct or more or less united; stigmas elongated, acute, either filiform or somewhat conic tapering to the point; upper part of capsule separating from the persistent base & dissepiment by circumscission. Flowers mostly sessile and glomerate; calyx always free, not ruptured. Natives of the old world, a few species introduced into different parts of America.

1. Cuscuta Epilinum (Weihe): Stem nearly hairlike, flowers sessile in sessile rather distant heads, 5-parted; segments of calyx broadly ovate acute or abruptly acuminate, nearly including the globose corolla; tube of corolla as long as its ovate acute lobes inconnived lobes, stamens half as long as the limb; substaminal scales short, broad crenulate, rather square, at the truncate apex; styles half as long as the inconnate stigmas both of the together shorter than the length of the globose ovary. capsule globose, surrounded by the withered corolla. — Weihe Flora, Ann. des Apoth. 8. p. 51. Reichenb. ic. t. 693 Koch, syn. p. 495. DC prod 9. 452. C. densiflora Soy Fr. Chois Epilinella cuscutoides, Pfeiff. Bot. Zeit. 4. p. 20 Cuscuta europaea ? Hayne, Darlingt. fl. cestr 2d ed. p. 167 Hygrocrocis Parasitus in flax

Common in flaxfields in central & western Europe, and introduced with flaxseed into the eastern States, Pennsylvania, Darlington New York Beck Massachusetts, Oakes! In wet seasons this species becomes very destructive to flaxfields. The thin, nearly hairlike stem is mostly simple, nearly hairlike, pale, from one to two feet high; flowers whitish, very densely clustered, whitish, small; capsule large in proportion. In wet seasons this species becomes very destructive to the flax-fields, in which it is abundant.

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved MISSOURI BOTANICAL GARDEN

2. Lepidanche Engler in Pflzm Fam. Oxtob 1842
(with character extended) Cuscuta Linn and authors ~~ed~~ partly exclusive
Engelmannia Pfeiff in Bot Zeit. 3-4 673 18 (not of Torr. & Gray)

~~Styles two~~, 2 ~~chips~~ Styles two distinct or rarely united, frequently supported by a fleshy stylopodium; stigmas capitate; capsule ~~bursting~~, not opening, finally separating entire from the persisting calyx; flowers mostly pedicelled, cymosely umbellate or paniculate; calyx free or surrounded by a number of bracts which resemble the sepals. — Natives of America, Polynesia and eastern Asia, only one species is known to be in Europe, undoubtedly introduced.

§. 1. Styles united

1. L. gamostyla: stem very thick, climbing high; glomerules few flowered; ~~pedicels~~ flowers 5-parted; sepals free, imbricate, orbicular, nearly including the much smaller cylindrical corolla; lobes of corolla much shorter than tube, obtuse; anthers ovate nearly sessile, subterminal; scales mostly in the shape of two distinct dentated wings on the sides of the adnate filaments; ovary globose, styles united stigma 2-headed; capsule large covered by the withered marcescent corolla Linth. flor. tex. exsicc. ~~flor. tex.~~ III fr. ined.

Along the upper Guadaloupe, Cibolo and other streams in western Texas

F. Lindheimeri climbing over the bushes and small trees, 10 to 15 or 20 feet high; F. Lindheimer! August to October — This largest north american species corresponds in a curious way with the largest european species the C. monogyna; in both the stem is of the size of a middle sized Gorse quill; both climb on trees; both have a corymbed small flower, but large fruit; and both are the only ones with one style in their countries. The resemblance may be carried out still further and still by the structure of the capsule and the shape of the stigma both must be referred to distinct genera.

§. 2. Styles distinct

* Depressae: ovary and capsules depressed-globose, without a stylopodium; capsule membranaceous, thin, mostly of a light yellowish colour.

† withered corolla covering the top of the capsule

2. L. teniflora: stem threadlike; flowers mostly 4-parted, with short pedicells,

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5375

forming dense cymes; tube of the corolla cylindric twice the length of its the obtuse erect or spreading lobes, and of the ovate obtuse segments of the calyx; stamens shorter than the limb; scales ovate, coarse, shorter than the tube; styles as long as the (depressed) ovary (; capsule covered by the remains of the corolla).— Cuscuta Cephalanthi Engelm in Sillim. Journ. 43 pag. 336. t. 6 f. 1-6.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

On the margin of ponds and swamps near St Louis, Mo, ! also Beardstown Ill, Mr Geyer! mostly on herbaceous plants, principally Compositae, Vernonia, Aster, Eupatorium, but also on many others, which come within its reach, as Polygonum, Lysimachia, Lycopus, Saururus, etc; rarely on shrubby plants, as Cephalanthus, Salix or Cornus. July to October. — This species which is common about St Louis is easily distinguished by its small flowers (which are mostly 4- (rarely 3 or 5-parted) and the hoodlike remains of the corolla on the capsule; the cylindrical tube of the corolla becomes convolute as soon as the ovary enlarges, and finally tears loose from its base. — I have first described this species first under the name of Cuscuta Cephalanthi, under the mistaken idea that most Cuscutae adhere principally to one species or genus of plants, and only accidentally stray on others; but though this may be true as to some species to some extent, yet it generally is not so; most of them grow promiscuously on all the plants in their neighborhood, and however different in their structure, & their properties. In this especial case, I have found that Lepid. teneriflorum grows more rarely on Cephalanthus, than on most other plants already named — I have therefore found myself obliged to change all these names, derived from the plants on which these parasitics vegetate.

3. L. cuspidata (Engelm. sub Cuscuta): stem threadlike, much branched; flowers 5-parted, pedicelled, forming loose bracteose cymes; tube of the corolla cylindrical sepals free, ovate (the exterior cuspidate) imbricate, surrounded by similar bracts; tube of the corolla cylindric or dilated somewhat funnelshaped, longer than the calyx, and its ovate or convolute-ovate acutish erect or spreading lobes, and than the calyx; stamens shorter than the limb. scales ovate, fimbriate, converging reaching to the nearly throat of the corolla, connected with one another at base; styles filiform, several times longer than the globose ovary; globose capsule covered by the remains of the corolla. — Cuscuta

2. Cass. for Flora

cuspidata Engelm. in Boston Journ. N. Hist. V p — Planta Lindheim. I, 125

α. protensis: flowers smaller, fewer bracts around the calyx, tube of corolla more cylindric and not so large in proportion; stamens only half as long as the ovate acutish lobes of the corolla: styles twice 3 or 4 times as long as the ovary

β. humida: flowers larger, of thinner texture Linth. Pl. Lindh. ex sched. I, 125 surrounded with many bracts, tube of corolla somewhat funnelshaped, larger in proportion; stamen somewhat shorter than the lanceolate acute lobes of the corolla; styles 3 or 4 times as long as the ovary. — Linth. Pl. Lindh. ex sched. Bot. Lindh. II, 277.

Texas, from the Brazos to the Colorado, F. Lindheimer!

Var α in dry prairies west of the Brazos, in June & July, growing on Tephrosia, Bradburia, Ambrosia, Liatris etc. var β. in rich, moist bottom Lands on the Colorado, in August, growing on Iva, Ambrosia, Rudbeckia. — ~~Inflorescens loose~~ Pedicells with numerous cuspidate bracts, the uppermost surrounding the calyx like an involucrum, similar in shape but smaller than the sepals; sepals somewhat lacerate or crenulate ovate, carinate (carina less distinct in β.) cuspidate interior ones ~~cuspidate~~, rather obtuse, all convex, closely imbricated; lobes of corolla thin, membranaceous, with a strong (in β. a less distinct) midnerve, formed by large oblong or linear cells; when dry convolute; at the base the lobes are dilated and cover one another more than in any other N. am. species. Styles remarkably slender and long, vary very much in proportion, but soon after flowering ~~swelling~~ enlarging, and giving the corolla an urceolate shape.

†† bothered corolla remaining at the base of the capsule

4 L. arvensis (Beyrich in herb Berlin under Cassuta): stem filiform branched, flowers pedicellate, (small) 5-parted, pedicels unequal in loose often somewhat umbellate cymes, ~~calyx~~ deeply 5-parted ~~shorter~~ campanulate tube of the corolla, ~~smaller~~ included in or exceeding the deeply 5 spreading or reflexed parted calyx, and shorter than the lanceolate acuminate lobes of the limb, stamens much shorter than the limb, scales ovate, fimbriate converging ~~closing the tube~~, and ~~2~~ ovules exceeding ~~of~~ the tube; styles as long or more than the globose depressed 3 capsule globose surrounded at base by the remains of the corolla. —

more open α. laevis: calyx thin, membranaceous, either much wider than the tube of the corolla and then 5-angled at the commissures of the sepals, or ~~not~~ closely appressed, and then less distinctly angled, mostly longer, sometimes also shorter than the tube of the corolla ~~obtuse, open, rather open~~. — Cuscuta pentagona Engelm! S.Ill. in Journ 43, l.c. t.6 fig 22-24 & 45 p. 340 & p. 76
C. arvensis Beyrich! in ~~herb reg. Berol.~~ Berlin C. globularis Nutt! in herb Acad Phil.

β. verrucosa: calyx thick, verrucose, mostly much shorter than the deeply campanulate somewhat globose tube of the corolla. — Cuscuta verrucosa Engelm! S.Ill. Journ 43, p. 74. l.c. t.6 fig 25 a-e Linn. Pl. tex exsicc Linnh. I, 127 & III —

Virginia Rugel! N. Carolina Curtis! S. Carolina Beyrich! Bosc! Florida Rugel! var. α, with a broad angular calyx, in fields, pastures, dry places, growing on many Compositae (Aster, Erigeron, Chrysopsis, Pterocaulon, Ambrosia etc, also on Pamplona, Polygonum, frequently on Trifolium etc. A form with small calyx (C. pentagona β. minor calyx Engelm l.c. ~~t.6 fig 25~~) in ~~Illinois~~, Geyer! on sandy dry hillsides on Aster, Solidago, Artemisia, Malva, Euphorbia etc. A form with large but not angular calyx, and narrower lobes (C. pent. γ. calyptrata Engl. l.c.) in Texas, Lindheimer! on Solidago, Boltonia etc.

Var. β. in Texas Drummond! Lindheimer! from Houston to San Antonio, mostly on Petalostemon, but also on Tephrosia Yaupon, Dalea, Hedyotis, Chrysopsis, Solidago, Coreopsis, Brazoria, Ipagia, and even on Cornus. May to July. —, This is the smallest flowered of our ~~native~~ species and the earliest one; the shape and texture of the calyx are so different in the different forms of this plant, that I ~~had~~ distinguished two species till I ~~had~~ specimens collected by Mr Lindheimer near San Antonio variable in their calyx and ~~finding~~ finding some of the intermediate between the extreme forms, proved them to belong to one species; In all of the the calyx segments are obtuse ish to rounded, and the ~~segments~~ lobes of the corolla acute to long acuminate, ~~with~~ spreading or reflexed but the tips incurved; scales large, much fringed, ~~covering~~ (nearly the full ovary); styles mostly as long as the ovary, but ~~in~~ from San Antonio, They are longer.

5. L. umbellata (Humb. Bonpl. & Kunth? Torrey under Cuscuta): stem ~~hairlike~~ wiry; fascicles of flowers 5-parted, long pedunculate, forming few flowered umbelliform fascicles; segments of

the calyx ovate - lanceolate, acute, longer than the campanulate tube of the corolla; lobes of the corolla lanceolate, acute, twice as long as the tube, erect or spreading; stamens nearly as long as the tube, filaments ~~broad at base~~, subulate; ~~anthers linear oblong~~ styles ovate, finely extending to the throat of the corolla; ~~filiform~~ capillary styles about as long as the ovary. — Torrey in Ann Lyc. N.Y. 2, p. 223. Longs Exped. II. p. 86. Humb. B. & K. n. gen. et sp. 3. p. 121. - DC. prodr. 9. p. 460

Barren regions of the upper Canadian river, at the base of the Rocky mountains almost exclusively on Portulaca oleracea, Dr James! — Humboldts plant has been collected between Guanaxuato & Salamanca, on the plateau of Mexico, also on low prostrate plants, and the description so far as it goes appears to prove the identity of Humboldts & Torreys plant, but it was not possible to compare specimens, — in the mexican plant however the lobes of the corolla are reflexed, (in ours ~~erect~~ slightly spreading) and filaments linear, (in ours distinctly subulate ~~from a broad base~~). — This species is intermediate between L. arvensis & L. chlorocarpa but distinguished by the low growth the finer, harsher stems, the long pedicels, flowers much smaller than in L. chlorocarpa, lobes of calyx & corolla more elongated; styles ~~thinner than~~ slenderer, valves less divided, lobed, stamens shorter, anthers more elongated. The texture of all parts of the flower is very fine.

6. L. chlorocarpa: ~~stem~~ thread like, flowers mostly 4-parted, ~~sessile~~ on short pedicells, subsessile glomerate; tube of the corolla campanulate shallow nearly equal to its acute, spreading lobes, and the ~~acute~~ triangular segments of the calyx; stamens as long as the limb, scales (small) mostly bifid, inclosed, appressed; styles thick, as long as the depressed ovary; ~~remains of~~ withered corolla persistent at base of the depressed - globose capsule. — Cavanta Polygonum Englm! Ic. t. 6 fig. 26-29.

Wet places, margin of ponds etc Stlouis Mo, Drummond! Lindheimer! Englmann! — Along the upper Illinois rivers Englm! Mostly on Polygonum sp. but also on Panthorum, Lobelia, Aster, Ambrosia, Rudbeckia, Lycopus, Acalypha, also rarely in Prairie and on Cephalanthus.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

August to September. — Closely related to the two last ones; flowers much larger, more open, with shorter pedicels, segments of corolla shorter, ½ the length of the stamens, filaments & ovate scales smaller, ovary clavate, protruding out of the shallow tube of the corolla. Stems orange colored, flowers greenish white thin & membranaceous. — MISSOURI BOTANICAL GARDEN under GEORGE ENGELMANN PAPERS

7. L. californica (Choisy sub Cressula): stem hairlike, much branched; flowers 5-parted, pedicelled, in loose few-flowered fascicles; tube of the corolla campanulate, at the apex recurved, short, much shorter than the ovate lanceolate, acuminate, segments of the calyx; lobes of the corolla linear-lanceolate, acuminate, erect or slightly spreading, 3 or 4 times as long as the tube; stamens much shorter than the limb; anthers linear, equal to the filiform filaments; scales ? styles setaceous, 3 or four times as long as the globose ovary.

Upper California, Douglas, Sta Barbara Nuttall! or Sandwich Islands. — Choris. californica Choris Ces. p. 183 DC. prod 9, p. 457; Hook bot Beech suppl. p. 364. C. acuminata, Nuttall! in herb Lind Philad.

Upper Calif. Douglas Sta Barbara, Nuttall! or Sandwich Islands. — A very distinct species, with its few-flowered fascicles, elongated flowers, even the flowerbuds present the characteristic appearance both of the long pointed corolla, and the recurved points of the calyx. Calyx & corolla membranaceous very thin, tolerably scarious and semitransparent; scales have not been able to discover in the few imperfect flowers examined; styles very thin & slender, stigmas small, globose. — I have little doubt that Choisy's the plant of Douglas, described by Choisy, being the same as that of Nuttall described here; but he says that the under surface is but he mentions the styles as short and the stamens as included, hardly reaching over the tubular throat of the corolla.

** Umbonatae: ovary & capsule crowned with a stylopodium, ovary turnid, forming a fleshy stylopodium, which sometimes considerably elongated; capsule more or less umbonate or rostrate, of a firmer texture and mostly brown colour.

0 1 2 3 4 5 6 7 8 9 10 cm copyright reserved MISSOURI BOTANICAL GARDEN

3. Cuscut. for Flora sepels united into a tube

† flowers pedicelled heads few and distant, corolla mostly remaining at the base of the capsule.

8. C. Gronovii (Willd ~~var. Roth Sch. 6 p. 205~~ ; stem threadlike; flowers ~~mostly~~ 5-parted, (~~capitat~~ - var. ~~pedunculata~~) on shorter or longer pedunles, forming small glomerules, or smaller or larger, denser or looser panicles; tube of the corolla deeply globose-campanulate, longer than the pellucid-punctate, (creat or spreading (finally reflexed) lobes and the ovoidly obtuse more or less oval) sometimes carinate and ~~slight~~ finely crenulate lobe segments of the calyx; stamens as long or a little shorter than the ovoid scales fimbriate, convergent, mostly united at base; ~~ovary~~ styles ~~slender~~ as long or longer than the ovary; capsule corolla ~~remaining~~ withering mostly at base of the imbonate capsule. —

Cuscuta subuliflora etc Gron. Virg. 18 (for wilded) C. americana Linn. and mant. fl. amer. partly ~~Chois Cusc. p. 185 t. 4~~

v Cuscuta Gronovii Willd ~~in R. & Sch. 6. p. 205.~~

~~f. 3. DC. prod. 9 p. 459.~~ C. vulgivaga Engelm. ! l. c. p. 338 t. 6. f 12-16

~~α laxiflora : flowers larger in loose very much branched cymose panicles.~~

α. paviflora : flowers smaller, mostly on shorter pedicells forming umbells or even glomerules; — ~~C. Gronovii~~ withering corolla always at base of capsule. — Cuscuta C. Gronovii Herb Willd! n. 3160! Chois. Cusc. p. 185. t. 4. f. 3. DC. prod. 9. p. 459

C. Sanamosa Nutt! hby Acad Phil. β. laxiflora : pedicells longer flowers larger in loose very much branched cymose panicles; withering corolla at base of capsule.

γ. calycostata : pedicels more or less bracted, flowers larger, deeply campanulate (capsule) approaching to cylindrical ~~subcylindrical~~, tubulous; ~~stylescales~~ scales oblong linear, slightly carinate; ovary with a larger stipe-podium, styles ~~very slender~~ equalling much longer capsule ovoid, umbonate, surrounded, and often even covered by the withered corolla

δ. tetrameris : flowers in compound umbelliform cymes, 3-4-parted.

Throughout the United States, var α from Vermont! & New Hampshire in the eastern States to Georgia! β. mostly west of the Alleghanies, western New York! to Wisconsin! Missouri! Kentucky! & Alabama! also in Pennsylvania! & N Carolina! γ in Texas! δ in Connecticut!

On very different plants, dicotyledonous and monocotyledonous innocuous and perfonous herbaceous and shrubby; especially on many Compositae, (Have seen it on Vernonia Eupatorium Aster, Solidago, Bidens) on Polygonum, Desmodium, on Ulex & Robinia, on

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Vhus toxicodendron Cephalanthus, Sambucus, Cornus, Ribis, Laurus
Bignonia, Carpinus, Rubus, Salix etc. ++ ✱ August to October —

This most variable species may always be recognized by the
duply campanulate
characters given; the calyx is always shorter than the tube of
the corolla, its segments always broadly ovate, obtuse, and all or
inbricate
at least the outer ones carinate by a number of warts or glandular
protuberances; the lobes of the corolla are mostly only half
as long as the tube, sometimes of the same length, broadly ovate
obtuse; styles always slender, ~~sometimes~~ mostly longer than
the ovary; capsule, globose, umbonate or somewhat
ovoid. The northeastern form is ~~mostly~~ small flowered
with very short pedicells, and appears to be a small, low plant;
the southeastern has rather
longer pedicells (Wilddenow, specimen for example, which
will
does ~~not~~ agree with ~~Chamisso~~, his short diagnosis: floribus
paniculatis
~~corymbosis~~, but not at all with Chamisso's l.c.: floribus
arctissime glomeratis), the usual plant (often creeps over
high bushes, 5 to 8 feet high; is larger, with thicker stems,
and large loose decompound, rarely more dense panicles)
stem reddish orange, flowers white.

9. L. patens: stem threadlike; flowers 5-parted, pedunculate,
crowded in dense umbelliform or paniculate cymes; tube of the
corolla ~~short~~ campanulate, wide and shallow, equal in length
to the linear oblong or oblong obtuse spreading lobes, and ~~mostly~~ a
little longer than or equal to the ~~oval~~ oblong obtuse entire & smooth
segments lobes of the calyx; stamens generally as long as the limb;
filaments subulate; scales large, pinnatifid laciniate,
convergent, mostly exceeding the tube; styles thick, as
long as the ovary, withered corolla persistent at the base of
the globose, ~~very~~ slightly umbonate capsule. — Cuscuta Saumi,
Engelm. l.c. 43. p. 339 t. 6. f. 17–21 (pg. 18. bad) and 45 p. 14

β? glandulosa: flowers on short pedicels in glomerules,
shallow open tube of the ~~small~~ ~~glandulous~~ corolla about the
same length as the very glandulous calyx, and shorter than
lobes of the corolla; stamens ~~rather~~ a little shorter than the spreading
scales very large, styles thick as long as the glandulous
ovary; capsule very glandulous, depressed.

On the margin of ponds & swamps Missouri! Illinois!
to Kentucky, Short! and western New York! A Gray! locally on

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Polygonum, but also on Aster, Veronica, Eupatorium Purpureum & also
on Cephalanthus, Salix, & Alnus too. Septemb. & Octob.
β on Polygonum, fl. Florida, Rugel! to Louisiana, Sartwell!

— Very near C. Gronovii, stem also yellow or orange-coloured,
fully as thick but never climbing high
always distinguished by the very open flower, which is nearly
smallest than at least the western form of C. Gronovii,
but bears a larger, more depressed, less umbonate
capsule; and by the longer but narrower lobes of
calyx and corolla; which latter are not carinate, the
segments of the calyx are not carinate nor crenulate,
& the corolla is rarely and only slightly dotted. —
Var β. on the contrary is glandular all in most part
of the flower, and moreover so far as I could
ascertain in the dried specimens before me, without any
stylopodium on the ovary or elevation on the capsule;
Note. this however is not membranaceous as in the allied
species without a stylopodium, but coriaceous; it
may prove to be a distinct species.
Specimens, sent from N. Carolina? by Mr. Curtis & distinguished by much smaller
nearly 4-parted flowers may belong to some
variety of C. patulus

10. C. rostrata (Shuttleworth under Cuscuta): stem threadlike, thick;
flowers (large) 5-parted, pedunculate, in umbellate cymes; tube of the
corolla globose-campanulate, twice as long as the ovate obtuse
slightly crenulate segment of the calyx and the spreading
(finally reflexed) ovate obtuse lobes of the corolla; stamens
nearly as long as the limb, scales fimbriate, at base
united; styles filiform, slender, & about the length of the
ovary, with its equally long stylopodium; withered corolla
persistent at the base of the (very large) rostrate
capsule. — Cuscuta rostrata. Shuttleworth in Rugel's coll.
Exsicc. in Boston Journ. N. Hist. 5 p.
Shady with places
In the valleys and on the declivities of the Alleghany Mountains
from Virginia to South Carolina, Rugel! Buckley! Gray & Sullivant! Curtis! on Aster, Solidago, Rudbeckia, Eupatorium, S. Ophioides, also on Vitis & Rubus. August to
October. — Calyx & Corolla of the same shape and
proportions, as in C. Gronovii, but larger,
usually large corolla destitute of the large carrells with give
the corolla of that species a pellucid dotted appearance.
tissue of the corolla regularly of somewhat elongated, hexagular

cells, not of linear elongated ones; ovary ~~remarkably~~ remarkably
prismatically through the very pointed ovary as well as capsule
it is nearly (× as long as the ovary proper, though only half as thick) the stylopodium being more developed than in any other of our
species; flowers very different in size. some are not larger
than the larger forms of L. Gronovii, while others are full
twice as large, to the length of 3 lines; capsule very large,
fully 3 lines long, globose, attenuated to a stiff point.

11 L. connivens crenulata: Stem threadlike; flowers mostly 4-~~parted~~
(more rarely 5-) parted, pedunculate, in umbellate cymes;
tube of corolla cylindric, as long or a little longer than the ovate,
acute crenulate, erect at the apex inflexed lobes, and the
ovate acute carinate segments of the calyx; stamens
~~a little~~ somewhat shorter than the lobes; scales consisting of
a ~~few~~ ~~short~~ a few teeth at ~~both sides of the~~ adnate
part ~~of the filament~~; styles slender as long or a little longer
than the ovary with the very distant stylopodium; capsule
depressed, slightly umbonate, surrounded by or sometimes
covered with the remains of the corolla. — Cesati Consp.
Engelm! l. c. 43 p. 337. t. 6. f. 7-11. Cambrosa,
Beyrich! in herb Berlin, Hooker? in fl. Bor. am. 2. p. 77
C. parviflora Nutt. in herb Acad Phil.
Western Virginia, Gray & Sullivant! to Georgia Beyrich! and
Missouri! & Illinois! about St Louis mostly on Conyles, or on
such petioles, but mostly moist places, where hazels had
been growing; even on young hickories, also on Salix, on
Rhus toxicod; also on Eupatorium ~~ageratoides~~ Solidago Teucrium
Desmodium, etc; in Virginia on Ceanothus & Helianthus.
August to October. — This together with the next species
is distinguished by the remarkable structure of the corolla
from all others in our flora; it is fleshy, not membranaceous
with a somewhat even surface and an irregularly crenulate
margin, showing under a glass similar numberless nearly globular semi-
transparent cells. All subfilamentous scales smaller than
in any other N. american species sometimes even wanting
mostly consisting of one or two teeth on each side of the
adnate part of the filament. Flowers pale, flowers
white small, capsule large, brown. Flowers in the

53777

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

4 Carex. for Flora

Georgia and in some of the Virginia specimens as well as in most Missouri specimens 4 parted, in other specimens from Virginia and a few from Louisiana & Illinois 3 parted. Bayard's specimens of his C. umbrosa in the Berlin herbarium has 4 parted flowers and on short pedicels, with cylindric corolla, very distinct scales etc; — but his description in Hook l. c. leaves it very doubtful whether he could have had the same plant in view, as he there speaks of 5 parted sessile flowers, fimbriate scales etc.

12 C. porphyrostigma: Stem threadlike; flowers 5-parted, on shorter or longer, often very long peduncles, forming forming umbellate or paniculate, often very loose and few flowered cymes; tube of the corolla campanulate or turbinate-campanulate, as long or longer than the ovate or ovate lanceolate acute more or less carinate segment of the calyx, equal in length to or somewhat shorter than the ovate more or less acuminate, irregulate often one-nerved spreading at the apex, incurved lobes of the corolla; stamens equal to or shorter than the tube; filaments subulate; scales ovate lanceolate fimbriate, convergent, nearly equalling the tube; ovary with a very distinct stylopod, nearly equal in length to or shorter than the, nearly equalling the styles.

α. hispidula: flowers small, peduncles cymes loose few flowered, somewhat hispid, or nearly smooth; flowers on very long peduncles, small, segments of calyx short and long acuminate lobes of corolla longer than the tube. — Carex hispidula, Engelm! l.c. 45, p. 75. — Lindh. for. tex. exsicc. III —

β. neuropetala: cymes umbellate paniculate often many more compact, many flowered, nearly smooth, flowers larger, segments of calyx and the tube short acuminate 1-nerved lobes of corolla equal to the tube. Carex neuropetala, Engelm! l.c. 45. p. 75 Lindh. for. tex. exsicc. I. 123

γ. littoralis: cymes somewhat umbellate paniculate, flowers nearly smooth flowers very large, tube of the corolla wide & shallow, about equal in length segments of calyx broadly oval, acutish, hardly carinate, lobes of corolla also broad, abruptly acuminate, nearly 1-nerved, about equal in length to the wide & shallow tube of the corolla. — Carex neuropetala γ. littoralis, Engelm in Boston Journ. II. Hest 5 p

West Indies
Southern Illinois! Florida, Rugel! Upthe steppes of Kansas
Fremont! 2d Exped nro. 485 and common in Texas, Lindheimer! (B);
~~Var. β. the more common form in all these~~ found on
Compositae, Solidago, Helianthus, Rudbeckia, Liatris, also
Euphorbia Pctalostemon,
on Eysenhardtia, Myrica, and others; July to Septemb. Var. α.
on dry soil
in Texas from Houston to San Antonio Lindheimer! on
on Soledago, Ambrosia, Gaillardia Evolvulus, Mimosa, Eysenhardtia Abies
also on Commelyna; April and May. Var. γ. Galveston beach
of Galveston island, Texas Lindheimer! on Lyci— ranlcia
Borrichia, Iva, Salicornia; May. — ~~A well distinguished but~~
A very variable plant, but always easily recognized by the
above characters and the structure of the corolla, which entirely
resembles that of L. crenulata; anthers yellow as in all others,
or purple, stigmas always purple, deep brown when dry.
In the smaller, loosely flowered form, α. we frequently observe
a sort of pubescence of pellucid, vesicular hairs on the
pedicells and ofs also on the calyx, the only instance of this
in any N. American species. —

†† flowers sessile, surrounded by ~~bracts~~ imbricate
bracts, which in all respects resemble the distinct sepals; corolla
remaining on top of the capsule. Lepidanche Engl. (Ill. Jour)

13. L. comparta (Juss. mss. Choisy): stem threadlike, thick;
flowers densely glomerate, closely sessile; ~~glomerules~~
~~mostly often forming a~~ sepals subulate-fibrous ovate or orbicular obtuse,
crenulate, concave, closely appressed, imbricate surrounded by
2 to 5 similar but larger bracts; tube of the corolla
cylindric more or less exsert, twice as long as the 3-parted obtuse lobes;
stamens shorter than the limb; anthers
connate-pennaliform, convergent (united at base); ovary with a distinct
stylopodium equalling the styles; capsule globose, acutish,
~~mostly~~ 1 to 4-seeded. — Cuscuta comparta Juss. mss. in
Choisy Cusc. p. 185. t. 4 f. 2. DC. prod. 9. p. 458.
~~α. coronata~~: Cuscuta americana Mich. Fl. am. sept. 1. p.
α ant. fl. am. partly.

α. coronata: glomerules closer, more compact, often
in one long, thick, ropelike wreath; flowers smaller
tube of the corolla slender, only half included in the
calyx; capsule nearly globose, appearing long pointed by

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

the remains of the woodlike) of the corolla. — Cuscuta corniculata, Beyrich! in herb Berlin, C. compacta Choisy l.c. in part & Engelm.! in Boston Journ N. Hist 5. p.

B. adpressa: glomerules fewer flowered, not so close; flowers larger, tube of the corolla wider & shorter, scarcely nearly entirely included in the calyx; capsule acute appearing less pointed. — Lepidanche Cuscuta adpressa Engelm! Sill. Journ 45. p. 77.

Mostly on ligneous plants (in the mountains) from North Carolina, Gray & Sullivant! Curtis! to Georgia, Beyrich! & Alabama & Gates! On Corylus rostrata, Andromeda axillaris? Pinus glabra (α); β, near Philadelphia Dr Lauffinger! in Virginia, Rugel, Gray & Sullivant! and in Missouri & Illinois Engelm! on Ceplalanthus, Lonicera Diervilla, Cornus, Alnus, Ostrya, Salix Fraxinus, Bignonia, Vitis, (Smilax & Rhus Toxic.); more seldom on herbaceous plants. Polygonum, Eupatorium, Aster. August to Septemb. — Stems thick, glomerules of flowers very dense, forming masses of flowers several inches long and 1 or 2 inches thick; flowers greenish white, sometimes whitish, in S. generally turning brown in drying; the much longer exserted in α and giving the full grown capsule a pointed appearance, which is very striking.

14. L. glomerata (Choisy under Cuscuta): stem thread like, evanescent; flowers 5-parted, closely sessile, densely glomerate, forming long thick rope like coils; sepals membranaceous ovate or ovate lanceolate, obtuse, crenate, with squarrose reflexed tips, subulate surrounded by 5 to 10 similar (the exterior gradually smaller) bracts; tube of the corolla cylindrical or a little wider above than below somewhat exserted twice as long as the oblong obtuse spreading or reflexed lobes; stamens as long as the lobes; scales pinnatifidly laciniate, convergent, united at base; styles slender much longer than the ovary with the stylopodium; stylopodium nearly the size of the ovary proper or larger; capsule nearly globose, pointed, 1—2 seeded. — Cuscuta glomerata Choisy. Conc. p. 184 t. 4. f. 1. DC. prod. 9. p. 458. Lepidanche Compositarum Engelm! Sillim Journ 43 p. 343. t. 6 f. 30—35.

Wet prairies, Indiana, Dr Clapp! to Illinois & Missouri Drummond (Rich!) Engelm! Mostly on Compositae, Solidago

S. Ophion, Helianthus, Veronica; in Mr Richl's nursery it was very destructive to young pear trees. August & September

I summer the numerous orange red stems of this curious species appears, and sometimes over whole patches in low praires, so that they can be distinguished from afar. About the beginning of August the stems & many plant. Stems orange red, disappearing about the flowering time; that part only, which is closely appressed to the stem of the mother plant becomes flattened, swells out on both upper and under lower edge in irregular lines or knots; the epidermis soon bursts there and the little squarrose scaly bracted flowerbuds make their appearance; they grow rapidly, so that soon all the glomerules become merged in the thick (6–10 lines) ropelike coil of a white or pale straw coloured squarrose bracts & sepals; wherein the flowers appear only here & there — being almost hidden.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS
3378

Lepidanche tenuiflora (Cuscuta Cephalanthi Engle.)
on Cephalanthus, Vernonia, Eupatorium, Polygonum, Lysimachia
Lycopus, Teucrium, Bochmeria, Aster, Cornus, Salix
S. de Saururus

St Louis, Beardstown,

The only variation was found on a large Polygonum,
flowers a little larger, lobes of calyx larger
but not as large as the tube — stamens
substaminal & only large the usually.

Lepidanche cuspidata (Cuscuta cuspidata Engelm. in Bost
Journ. N. Hist. 5, p.)
on Liatris, Ambrosia, Bradburia, Tephrosia
β. on Iva, Ambrosia, Plucheа
Texas, Prairies between the Brazos & Colorado & in the
Colorado Bottom lands
two varieties vide R. Lindheimer,

Lepidanche arvensis (Cuscuta arvensis Beyrich ined!)
Cuscuta pentagona Engelm) Cuscuta globulosa Nuttl in
hb. Acad Phil) × Monarda? (H Gray & H Shuttleworth)
α. pentagona on Fragia (?) (Virginia) on Aster, Ambrosia, Triplasis pra
Engelm. rand & shrubs — Bignonia, Plantago
Polygonum Sonchus Paniflora Lepidium. (N. Carolina
Curtis, Beyrich Carolina Beyrich Hab. Berlin; Bosc. Herb. Neuf
chatel) — Florida Angel in Hall Shuttlew — on Pleuraulon
Chrysopsis × graminea

β. microcarpa × On Ceanothus, Malva, Aster, Solidago, × Tenesia
Euphorbia, Helianthemum, × Pleuris aquilina
& Graminea (on Pleuris aquilina & Equisetum
hyemale it did not root though it formed rootlets
but the papillae do not enter the epidermis —
but such was the case on some graminea)
Beardstown Ill.

γ. calycina On Boltonia diffusa & Solidago
April to June, Prairies west of Houston. Texas
This possibly also = Georgia (Boykin) fragmentary
1 specimen

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5379

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

2

1. Lepidanche chlorocarpa (Cuscuta Polygonorum E.)
a Polygonum, Anthemis, Lycopus, Aster, Cephalanthus,
Bidens (~~Gymnolomia~~ Liatris) Ambrosia, Gnaphalium
(slight impression) — Atalypha

St Louis. H Penn Illin, also Drummond — H.C. Shuttleworth

Lepidanche glandulosa ~~Florida~~ on Polygonum
Florida (Angel & H Shuttlew.) and Louisiana
(Tuscumbia H.C. Aug 28th)

Lepidanche imbellata ? (Humb. sub Cuscut.) Torrey :-
Lyc. N.Y.) -- base of the Rocky Mts about
the Canadian River ? (~~Iowa~~ James - H.C.
Torrey)

Lepidanche vermicosa (Cusc. vermicosa E.)
Texas (Drummond III, 247; Lindheim I. 127.)
a Petalostemon. —

P. near Brazos Texas (Lindh. III....) on
Gaura et vide MSS. Drummond

Lepidanche californica (C. calif. Choisy)
Upper California, on Eriogonum (Nutt. H.C. to Phil)
near Sonoma

Lepidanche Gronovii (C. Gronovii W. Choisy.
C. vulgivaga E. C. americana auct).
Cephalanthus, Rhus tox. Polygonum, Sambucus, Amphicarpa
Diospyros, Rubus, Lonicera, Desm. Cornus, Vitis
Eupatorium, Aster, Bidens, Solidago

Sambucus & vernonia Beardstown Geyer
Polygonum Kentucky, Short
Philadelphia Jungkegen, Lea
Ohio Frank on Hydrangea
Decatur New York, Gray
Georgia H.C. Carey
North Carolina Gray on Polygonum etc, Buckley
Alabama

5380

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

3

Lepidanthe Gronovii contd.

var. calyptrata = stylis filiformibus, demum elongatis = Alabama, Buckley — Aster Salix — Texas Drummond, Hb Hook; Lindheimer — Carpinus, Polygonum (on Rhus toxit) Cephalanthus, Aster, Veronica, Gramineae (stylus very long) Rochmeria, Rhus tox. Solidago Virga

var. minor. Georgia Boykin, Solidago, New hampsh. J. Carey — Vermont on trees in J. Carey — Salem. mass (Hb. Acad Ph.) — Solidago Boston (Hb Ac Ph)

= C. canescens mutt. Hb. Acad Phil.

Gronii Wills. herb. 3160

var tetramera = Hitchcock communicat H Carey

Lepidanthe patens (Cass. Saururus Eaton, probably only a variet of L. Gronovii

St Louis on Saururus, Polygonum, Penthorum, Cephalanthus, Aster, Veronica, Gramineae (without penetrating the Epidermis) Salix, Eupator, Atalypha, Hibiscus, Rhus tox Western New York = Saururus (A Gray)

Kentucky (Short)

var minor N. Carolina: Curtis

Lepidanthe reflexata for the Mountains of western Virginia to North & South Carolina

Ulmus Aster, Solidago Eupatorium, S. Gigantea Angel Buckley Gray & Sullivant — Curtis

Lepidanthe compacta & coronata (Cass. coronata Beyrich in Hb Willd. Berlin)

Georgia Beyrich — North Carolina Curtis Gray & Sullivant or Conyzoides reflexata Andromeda axillaris; Alabama Gates = hb. Gray on Prinos glaber

5381

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

4/ Lepidanche conjuncta

B. adpressa (Lepid adpressa Engl.

St Louis or Ohio Valley, Cephalanthus Cornus

Polygonum Fraxinus, Laurus Benzoin, Vitis

Eupatorium Asimina

hills on the Missouri or St Lix, Cornus, Alnus

Philadelphia (Dr Jaeger) - Portsmouth Virginia

Angel in Hb. Shuttleworth

Western Virginia or Ohio, Salix, Ohio?

A Gray & Sullivant

Lepidanche glomerata (Cassia glomerata Nees.

Lepid. composita Engl.)

Vermont Solidago S. Ohio Helianthus (yellow

pasture in the morning)

Missouri, Illinois & Indiana

Lepidanche ramulata (Cassia Carol. E, C. ambrosa

Asperifolia in Hb. Berlin (not Hook fl. bor.-amer.?)

Cassia marilandica Nutt in Hb. Berl. Oak Salix

St Louis on Carolus Carya

Aster oblongifol. Ohio Valley. Desmodium Verbena, Teucrium

Eupatorium Aster Solidago

Beaudette. - Solidago (Geyer)

Western Virginia - mountains near the warm Springs

on Ceanothus Helianthus

Georgia Asperifolia in Hb. Berlin

Cult on Euphorbia, Chrysanth. indicum Aster ericoides

Graminea (actually rooting) Amorpha, Verbena

Portulaca

Lepidanche porphyrostigma

α. hispidula Houston Texas, on Solidago, Aster,

Eysenhardtia, Mimosa - low - shrub

Neobraunfels April, Gaura, Mimosaceae (principally)

Gillaspie, Dallas, Commelyna - Rhus -

on Dry prairies, April.

β. neuropetala Illinois - Dry swamps near Kaskaskia

on Compositae.

Florida Angel in herb Shuttleworth on Compositae & Helianthemum

0 1 2 3 4 5 6 7 8 9 10 cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5382

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN